

# Culture Guide

## Genus

	Optimal Planting Time		Retail Sales Window			Moisture in Production			Requires Shade in Production	Vernalization Required for Bloom	Vernalization Beneficial
	Late Summer-Fall	Spring	Spring	Summer	Fall	Low	Average	Consistent			
Achillea											
Acanthus											
Agastache											
Ajuga											
Alchemilla											
Allium											
Amsonia											
Anemone (Japanese)											
Anemone <i>sylvestris</i>											
Antirrhinum											
Aralia											
Asarum											
Asclepias											
Aster											
Bergenia											
Brunnera											
Calamintha											
Callirhoe											
Campanula											
Caryopteris											
Centranthus											
Ceratostigma											
Chelone											
Coreopsis											
<i>Dianthus carthusianorum</i>											
<i>Dianthus gratianopolitanus</i>											
<i>Dianthus barbatus</i>											
Echinacea											
Eupatorium											
Galium											
Gaura											
Gentiana											
Geranium											

\*Many perennials, including spring bloomers, perform better when planted in late summer/fall prior to allow time for bulking and root development, but may be grown in spring as well.

\*All plants require consistent moisture until they begin to root into the container. Moisture in production relates to what the plants typically need after rooting is sufficient.

# Genus

	Optimal Planting Time		Retail Sales Window			Moisture in Production			Requires Shade in Production	Vernalization Required for Bloom	Vernalization Beneficial
	Late Summer-Fall	Spring	Spring	Summer	Fall	Low	Average	Consistent			
Grass-Bouteloua											
Grass-Calamagrostis											
Grass-Carex <i>flacca</i> , <i>C. oshimensis</i> , <i>C. pensylvanica</i>											
Grass-Carex <i>morrowii</i>											
Grass-Chasmanthium											
Grass-Deschampsia											
Grass-Festuca											
Grass-Hakonechloa											
Grass-Luzula											
Grass-Ophiopogon											
Grass-Panicum											
Grass-Pennisetum											
Grass-Schizachyrium											
Grass-Sesleria											
Grass-Sporobolus											
Helenium											
Heliopsis											
Heuchera											
Heucherella											
Hydrangea <i>arborescens</i>											
Hypericum											
Iris <i>cristata</i>											
Kalimeris											
Knautia											
Lamium											
Lavandula											
Leontopodium											
Lespedeza											
Leucanthemum											
Liatris <i>ligulistylis</i>											
Liatris <i>microcephala</i>											
Liatris <i>pycnostachya</i>											
Limonium											

# Genus

	Optimal Planting Time		Retail Sales Window			Moisture in Production			Requires Shade in Production	Vernalization Required for Bloom	Vernalization Beneficial
	Late Summer-Fall	Spring	Spring	Summer	Fall	Low	Average	Consistent			
Monarda											
Nepeta											
Origanum											
Penstemon											
Perovskia											
Persicaria											
Phlomis											
Phlox hybrids											
Phlox <i>divaricata</i>											
Phlox <i>paniculata</i>											
Phlox x <i>procumbens</i>											
Physostegia											
Polemonium											
Rudbeckia											
Sagina											
Salvia <i>greggii</i>											
Salvia <i>nemorosa</i>											
x Sedoro											
Sedum											
Senecio											
Sisyrinchium											
Solidago											
Stachys											
Stachys <i>byzantina</i>											
Tanacetum											
Thymus											
Tiarella											
Tradescantia											
Tricyrtis											
Verbena											
Vernonia											
Veronica (groundcover types)											
Veronica (upright types)											